

Peri-Operative Management of Gynecology Patient During COVID-19 Pandemic			
	Confirmed COVID or Suspect		COVID Screen Negative AND Asymptomatic
Anesthetic Technique	GA as medically indicated	Regional/Sedation preferred	Regional/Sedation General Anesthesia (GA)
OR Precautions	Airborne	Droplet/Contact	Standard OR Precautions
OR Location	OR C	OR C	OR A or B
In Room Principles	<ul style="list-style-type: none"> OR doors to remain closed Have runner outside (N95) 	Standard OR Practice	Standard OR Practice
Recovery Principles	<ul style="list-style-type: none"> Only Anesthesia & AA or RN to remain in OR for extubation Recover for 20mins* in OR, doors closed Transfer to Isolation room or 1P36 to cohort Runner remains available 	Direct to Isolation Room or 1P36 to cohort	Direct to PACU
Room Cleaning	Terminal Clean 20min** after extubation Droplet/Contact	Terminal Clean Droplet/Contact	Regular Clean

AIRBORNE: Fit tested N95 mask, plus droplet/contact: eye protection, surgical hat, gown, gloves

DROPLET/CONTACT: Surgical mask, eye protection, surgical hat, gown, gloves

*Based on local HVAC functionality, 20mins is the required room closure interval to clear aerosols in OR

Version History

DATE	DOCUMENT NUMBER and TITLE	ACTION TAKEN
29-Apr-2020	C-06-01-60631 COVID-19: Peri-Operative Management of Gynecology Patient	Developed by BCW Maternal Newborn Program; Approved by Professional Practice Director

Disclaimer

This document is intended for use within BC Children's and BC Women's Hospitals only. Any other use or reliance is at your sole risk. The content does not constitute and is not in substitution of professional medical advice. Provincial Health Services Authority (PHSA) assumes no liability arising from use or reliance on this document. This document is protected by copyright and may only be reprinted in whole or in part with the prior written approval of PHSA.